

Endothelial dysfunction and reduced heart rate variability in patients with metabolic syndrome

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Copyright E.N. Smirnova et al. According to experts of the World Health Organization (WHO), metabolic syndrome (MS) can be considered as pandemy of the XXI century, because its prevalence among the population of developed countries is about 25-35%. In this study with the purpose of complex investigation of the autonomic nervous system and endothelial function we included 66 patients with MS between the ages of 25 and 61 (46.9 ± 9.9 years). A comparison group of apparently healthy individuals (16 individuals, average age of 45.3 ± 2.3 years; $P > 0.05$) was studied. To evaluate the response of microvascular tone, we used the method of wavelet analysis of skin temperature oscillations during cooling of the limb. All patients underwent the study of heart rate variability. The levels of insulin, endothelin-1, and vascular endothelial growth factor were determined using enzyme immunoassay. Patients with MS had significant differences in all metabolic parameters. Our study showed that in the group of MS there is a decrease of the variability of heart rhythm compared with the healthy group. Conducting cold test revealed signs of endothelial dysfunction in the MS group, which was manifested by the decrease of the index of vasodilation in the endothelial and neurogenic frequency range. In the study group we determined the increase in biochemical markers of endothelial dysfunction, which correlated with parameters of vasodilation. Also, the presence of endothelial dysfunction significantly correlated with signs of reduction of the variability of the heart rhythm.

<http://dx.doi.org/10.4081/itjm.2018.847>

Keywords

Cold test, Endothelial dysfunction, Heart rate variability, Metabolic syndrome

References

- [1] Gurgonian SV, Vatinian SKh, Zelveian PA. Arterial hypertension in metabolic syndrome: pathophysiological aspects. *Ter Arkh* 2014;86:128-32.
- [2] Saladini F, Di Marco A, Palatini P. Autonomic dysfunction: how to identify and when to treat? *High Blood Press Cardiovasc Prev* 2016 [Epub ahead of print].
- [3] Bruno RM, Ghiadoni L, Seravalle G, et al. Sympathetic regulation of vascular function in health and disease. *Front Physiol* 2012;3:284.
- [4] Rajendran P, Rengarajan T, Thangavel J, et al. The vascular endothelium and human diseases. *Int J Biol Sci* 2013;9:1057-69.
- [5] Smirnova E, Podtaev S, Mizeva I, Loran E. Assessment of endothelial dysfunction in patients with impaired glucose tolerance during a cold pressor test. *Diabet Vasc Dis Res* 2013;8:1-9.
- [6] Cubbon RM, Kahn MB, Wheatcroft SB. Effects of insulin resistance on endothelial progenitor cells and vascular repair. *Clin Sci* 2009;117:173-90. nl

- [7] Favero G, Paganelli C, Buffoli B, et al. Endothelium and its alterations in cardiovascular diseases: life style. *Intervent Biomed Res Int* 2014;801896. e
- [8] do Carmo JM, da Silva AA, Wang Z, et al. Obesity-induced hypertension: brain signaling pathways. *Curr Hypertens Rep* 2016;18:58. l u
- [9] Barton M, Baretella O, Meyer MR. Obesity and risk of vascular disease: importance of endothelium-dependent vasoconstriction. *Br J Pharmacol* 2012;165:591-602. ia
- [10] Lobato NS, Filgueira FP, Akamine EH, et al. Mechanisms of endothelial dysfunction in obesity-associated hypertension. *Braz J Med Biol Res* 2012;45:392-400.
- [11] Franz R, Maturana MA, Magalhães JA, et al. Central adiposity and decreased heart rate variability in post-menopause: a cross-sectional study. *Climacteric* 2013; 16:576-83.
- [12] Wu HT, Hsu PC, Sun CK, et al. Assessment of autonomic dysfunction in patients with type 2 diabetes using reactive hyperemia. *J Theor Biol* 2013;330:9-17.
- [13] Indumathy J, Pal GK, Pal P, et al. Association of sympa-thovagal imbalance with obesity indices, and abnormal metabolic biomarkers and cardiovascular parameters. *Obes Res Clin Pract* 2015;9:55-66.
- [14] Gaudreault V, Després JP, Rhéaume C, et al. Exercise-induced exaggerated blood pressure response in men with the metabolic syndrome: the role of the autonomous nervous system. *Blood Press Monit* 2013; 18:252-8.
- [15] Campbell B, Aguilar M, Bhuket T, et al. Females, Hispanics and older individuals are at greatest risk of developing metabolic syndrome in the U.S. *Diabetes Metab Syndr* 2016 [Epub ahead of print].
- [16] Patel TP, Rawal K, Bagchi AK, et al. Insulin resistance: an additional risk factor in the pathogenesis of cardiovascular disease in type 2 diabetes. *Heart Fail Rev* 2016;21:11-23.